T is



wherein

R^1 is C_1-C_{10} alkyl;

R^3 is —C(O)C_1-C_5 alkyl, —C(O)N(H)C_1-C_{10} alkyl, or —(C_1-C_{10} alkylene)-$NR^{3a}R^{3b}$,

wherein R^{3a} and R^{3b} are independently in each instance, hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, and acyl, wherein alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, and acyl are optionally substituted;

R^4 and R^5 are, independently in each instance, hydrogen or C_1-C_5 alkyl;

R^6 is —OH, —O—, —$NHNH_2$, or —NHNH—;

R^7 is, independently in each instance, hydrogen, —OH, —O—, halogen, or —$NR^{7a}R^{7b}$,

wherein R^{7a} and R^{7b} are independently in each instance, a bond, hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, acyl, and amino acid residue, wherein alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, and acyl are optionally substituted;

R^8 is, independently in each instance, hydrogen, deuterium, —NHR^9, or halogen,

wherein R^9 is hydrogen, —C_1-C_5 alkyl, or —C(O)C_1-C_5 alkyl; and

m is one or two;

Q is —CH_2— or —O— wherein

when Q is —O—, then R^2 is C_1-C_{10} alkyl, C_1-C_{10} alkylene, C_1-C_{10} alkynyl, C_1-C_{10} alkynylene, a regioisomeric C_1-C_{10} triazolylene, —C_1-C_{10} alkylene-(5-membered heteroaryl), a regioisomeric —C_1-C_{10} alkylene-(5-membered heteroarylene), —C_1-C_3 alkylene-Q^1-$(CH_2)_n$aryl, —C_1-C_3 alkylene-Q^1-$(CH_2)_n$arylene, C_1-C_3 hydroxyalkyl, or C_1-C_{10} alkylether; or

when Q is —CH_2—, then R^2 is C_5-C_{10} alkyl, C_5-C_{10} alkylene, C_1-C_{10} alkynyl, C_1-C_{10} alkynylene, a regioisomeric C_1-C_{10} triazolylene, —C_1-C_{10} alkylene-(5-membered heteroaryl), a regioisomeric —C_1-C_{10} alkylene-(5-membered heteroarylene), —C_1-C_3 alkylene-Q^1-$(CH_2)_n$aryl, —C_1-C_3 alkylene-Q^1-$(CH_2)_n$arylene, C_1-C_3 hydroxyalkyl, or C_1-C_{10} alkylether; and

Q^1 is —CH_2— or —O—;

wherein said regioisomeric triazolylene is unsubstituted or substituted with alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, and acyl;

wherein said heteroaryl or regioisomeric heteroarylene is unsubstituted or substituted with alkyl, aminoalkyl, -alkylene-NH—, hydroxylalkyl, -alkylene-O—, carboxyalkyl, -alkylene-COO—, benzyl, or phenyl;

wherein said aryl is unsubstituted or substituted with nitro, amino, or —NH—; and

wherein n is an integer from one to five; and

k is an integer from one to thirty.

66. The compound of claim **65**, having a Formula A, B, C, or D: